

Laparoscopic Radiofrequency Ablation of Fibroids

Uterine fibroids (also called myomas) are non-cancerous growths that originate from the uterus. Up to 70% of women will have fibroids by age 50, and 25% of these women will have significant symptoms. Symptoms of uterine fibroids can include heavy menstrual bleeding, prolonged periods, pelvic pain, pressure on other organs (called bulk symptoms), or planning your life around periods. Symptoms often depend on the size and location of the fibroids in the uterus.

Laparoscopic radiofrequency ablation of fibroids (known as Acessa) is a procedure that uses heat to destroy fibroid tissue. Your surgeon places a camera through two small surgical cuts: one in the belly button and one in the skin over the lower abdomen (belly). Your surgeon looks inside your abdomen and inserts small instruments to destroy the fibroids (or growths) with heat and energy.

ABOUT THIS PROCEDURE


Why is this procedure performed?

Laparoscopic radiofrequency ablation of fibroids is used for women with symptomatic fibroids who want to save their uterus.


Benefits of the surgery:

Alternatives to this procedure can include:

- Saves the uterus
- Not having any treatment
- No surgical cuts on the uterus
- Medication to manage bleeding
- Faster recovery than traditional abdominal surgery done through a large incision
- Procedure to decrease blood supply to the uterus (uterine artery embolization)
- Removal of fibroids (myomectomy)
- Removal of uterus (hysterectomy)



Heat and energy destroying fibroid tissue inside uterus.



Laparoscopic method used to see and burn fibroids.

RISKS

What are the risks of this procedure?

There is a small chance of having a complication or problem when you have surgery. Your risk could be higher if you have had surgery before or have other medical conditions. Some risks include:

- Bleeding during or after the surgery
- Infection
- Injury to your bladder, intestines, or other structures near your uterus
- Blood clots in your legs or lungs
- Hernia (weakness and a tear in the wall of your abdomen)
- Need to switch to a laparotomy (surgery through one large surgical cut)
- Up to a 30% chance of recurrence of fibroids requiring further treatment or repeat surgery
- Complications in pregnancy including requiring a C-section for delivery and need for closer monitoring in pregnancy (please consult an obstetrician early in pregnancy)



A cross-sectional view of a laparoscope (camera) inside the abdomen

AFTER SURGERY

What should I expect after surgery?

Everyone recovers at a different pace after surgery. Many patients are able to get back to most of their usual activities by one to two weeks after surgery. Common symptoms after laparoscopic treatment of fibroids include:

- Irregular vaginal bleeding
- Fatigue or sleepiness from anesthesia
- Pain or cramping in your stomach and soreness from your surgical cuts
- Shoulder pain
- Constipation (difficulty emptying bowels)

Do not put anything in your vagina for at least two weeks (no tampons or sexual intercourse).

When should I call my doctor?

Serious problems after laparoscopy are uncommon, but notify your surgeon if you develop:

Difficulty breathing or shortness of breath

Heavy vaginal bleeding

Pain not controlled by your pain medications

Severe nausea and vomiting

A temperature over 100 degrees

Trouble urinating or having a bowel movement

Find an AAGL-trained surgeon near you

Search our directory of minimally invasive gynecologic surgeons

gynawareness.com